

GEYMBERG, V.G.; MEYER YEVA, N.P.

Investigation of properties of different variants of *Shigella sonnei*.
Zhur. mikrobiol. epid. i immun. no.11:49-54. N '54. (MLRA 8:1)

1. Is mikrobiologicheskoy laboratorii (zav. prof. V.N.Azbelev)
Otdela pishchevoy eieliyeny Instituta pitaniya AMN SSSR (Moskva)
(SHIGELLA,
sonnei, properties of different variations)

ASHBELEV, V.N.; BUDAGYAN, F.Ye.; GEIMBERG, V.G.

"Food poisonings and their prevention." M.I.Orlov. Reviewed by
V.N.Ashbelev, F.E.Budagian, V.G.Geimberg. Vop.pit. 13 no.1:59-61
Ja-F '54. (MLRA 7:1)

(Food poisoning) (Orlov, M.I.)

GEYMBERG, V G

1. The first part of the report is devoted to a description of the work done in the field of the theory of the structure of the atomic nucleus. The author discusses the results of the experiments carried out in the field of the study of the structure of the atomic nucleus, and also the results of the theoretical calculations. The author also discusses the results of the experiments carried out in the field of the study of the structure of the atomic nucleus, and also the results of the theoretical calculations.

GEYMBERG, V.G.

~~REDACTED~~

"Food-borne infection and intoxication." F.W.Tanner, L.P.Tanner.

Reviewed by V.G.Geimberg. Vop. pit. 13 no.6:51-53 N.D. 54.

(FOOD ADULTERATION AND INSPECTION)

(MLRA 8:1)

(TANNER, F.W.) (TANNER, L.P.)

AZBELEV, V.N.; GEYMBERG, V.G.; NEFED'YEVA, N.P.; RUBINSHTEYN, Yu.I.

"Sanitation bacteriology." V.I.Tets. Reviewed by V.N.Azbelev and
others. Vop.pit. 14 no.2:57-60 Mr-Apr '55. (MLRA 8:6)

(FOOD)

(BACTERIOLOGY)

(TETS, V.I.)

GEYMBERG, V.G.

Nutritional role and requirements of proteins. Vop.pit. 1/4 no.5:
45-47 S-O '55. (MLRA 8:11)

1. Otchet o diskussii, proiskhodiavshey na zasedanii Uchenogo
soveta v Institute pitaniya AMN SSSR.
(PROTEINS,
nutritional role)

GEINBERG, V.G.; NEFED'YEVA, N.P.
~~CONFIDENTIAL~~

"Dietetic and therapeutic sour milk products and the organization of
their production." A.M.Skorodumova. Reviewed by V.G.Geinberg, N.P.
Nefed'eva. Vop.pit. 15 no.5:58-59 8-0 '56. (MLRA 9:11)

(DAIRY PRODUCTS)

(SKORODUMOVA, A.M.)

GEYMBERG, V. G.; NEFED'YEVA, N. P.; RUBINSHTEYN, Yu. I.

"Microbiological examination and sanitary inspection of food
stuffs." S. P. Askalonov, I.B. Dobrier, B. L. Gordin, Reviewed
by V. G. Geinberg, N. P. Nefed'eva, IU. I. Rubinshtein. Mikro-
biologiya 25 no.2:243-245 Mr-Apr '56. (MLRA 9:7
(FOOD ADULTERATION AND INSPECTION)

CHYMBERG, V.G.

Tenth session of the Institute of Nutrition of the Academy of
Medical Sciences of the U.S.S.R. Vest.khir. 77 no.11:50-61 N '56.
(NUTRITION) (MLRA 10:1)

GEYMBERG, V. G.

YUGOSLAVIA / Microbiology. Antibiosis and Symbiosis.
Antibiotics.

F-2

Abs Jour : Ref Zhur - Biol., No 8, 1958, No 33788

Author : Geymberg, V. G.

* Inst : Not given

Title : The Significance to the Organism of Normal Intestinal
Microflora.

Orig Pub : Vopr. pitaniya, 1957, 16, No 5, 44-52

Abstract : No abstract.

1. The significance of the normal intestinal microflora to the organism of the rat (Rattus norvegicus) is studied. It is shown that the normal intestinal microflora is necessary for the development of the organism and for the maintenance of its health.

Card 1/1

GEYMBERG, V. G.

USSR / Microbiology. Microorganisms Pathogenic to Humans and Animals.

P-5

Abs Jour : Ref Zhur - Biol., No 20, 1958, No. 20930

Author : Geymberg, V. G.; Mikhlin, S. Ya.; Pavlova, Z. M.

Inst : Not given

Title : The Problem of Suppression of Enteric Microflora

Orig Pub : Zh. mikrobiol., epidemiol. i immunobiol., 1958, No 3, 128-129

Abstract : No abstract given

Card 1/1

GEYMBERG, V.G.
MIKHLIN, S. Ya., GEYMBERG, V.G., PAVLOVA, Z.M.

The role of microflora in the destruction of enterokinase in the large intestine of the rabbit [with summary in English].
Vop.mod.khim. 4 no.1:8-14 Ja-F'58 (MIRA 11:5)

1. Laboratoriya fiziologii pishchevareniya i laboratoriya mikrobiologii Instituta pitaniya AMN SSSR, Moskva.

(PROTEASES, metabolism
enterokinase destruction in large intestine in rabbit
role of intestinal bact. (Rus))

(PHOSPHATASES, metabolism
destruction in large intestine in rabbit, role of
intestinal bact. (Rus))

(INTESTINE, LARGE, bacteriology
bact. role in destruction of enterokinase & phosphatase
in rabbits (Rus))

GEYMBERG, V.G.; MIKHILIN, S.Ya.; PAVLOVA, Z.M.

Inhibiting normal enteric microflora. Vop.pit. 17 no.1:92-93
Ja-F '58. (MIRA 11:4)

1. Iz laboratorii fiziologii pishchevareniya (zav. - prof. G.K.
Shlygin) i otdela pishchevoy gigiyeny (zav. - prof. F.Ye.Budagyan)
Instituta pitaniya AMN SSSR, Moskva.
(INTESTINES--BACTERIOLOGY)

SHTENBERG, Abram Il'ich; PLOTNIKOVA, Yuliya Il'inichna; MUKHORINA,
Klavdiya Vasil'yevna; Primalni uchastiye: GEYMBERG, V.G.;
NEFED'YEVA, N.P.; NOVIKOV, Yu.V.; NATANSON, A.O., red.;
BUL'DYAYEV, N.A., tekhn. red.

[Guide to practical work in nutritional hygiene] Rukovodstvo k
prakticheskim zaniatiyam po gigeno pitaniia. Moskva, Medgiz,
1961. 358 p. (MIRA 15:7)

(NUTRITION)

RAZUMOV, M. I.; GEYMBERG, V. G. (Moskva)

Symbiosis of reticular cells with Bacteroides in the lymphoid
follicles of the intestine of rabbits. Arkh. pat. no. 4:24-31
'62. (MIRA 15:4)

1. Iz laboratorii patologicheskoy morfologii (zav. - doktor
meditsinskikh nauk M. I. Razumov) i bakteriologicheskoy laboratorii
Instituta pitaniya AMN SSSR.

(INTESTINES--MICROBIOLOGY) (BACTEROIDES)
(LYMPHOID TISSUE--MICROBIOLOGY)

NOSKOVA, Glafira Leonidovna; PEK, Georgiy Yul'yevich; GEYMELIN, V.G.,
kand. biol. nauk, nauchnyy red.; KAPLUN, M.S., red.;
BRODSKIY, M.P., tekhn. red.

[Effect of vacuum on the microflora of refrigerated food
products] Vliianie vakuuma na mikrofloru okhlazhdennykh pi-
shchevykh produktov; nauchnye soobshchenia. Moskva, Gos-
torgizdat, 1963. 37 p. (MIRA 16:7)
(Refrigeration and refrigerating machines)

Geymos, K. A.

USSR / Farm Animals. Cattle.

Q-2

Abs Jour: Ref Zhur-Biol., No 12, 1958, 54757.

Author : Geymos, K. A.

Inst : Not given.

Title : On the Causes of the Poor Reproduction of a
Herd (from the Experience in the Rearing of
Young Cattle).

Orig Pub: Zhivotnovodstvo, 1957, No 5, 73-76.

Abstract: The author analyzes his long-standing experience as well as certain experimental data, in regard to rearing calves. He reaches the conclusion that a full-value and steady feeding of the offspring and of the mother-cows constitutes the basic condition for the preservation and adequate development of young cattle. He considers the evaluation of the local feedstuffs to be of great importance.

Card 1/1

24

TOKAREVA, B.A., kand.med.nauk; GEYMOS, Ye.K.

Use of corticosteroids in the treatment of sympathetic ophthalmia.
Vest.oft. no.3:29-31 '61. (MIRA 14:9)

1. Glavnoye otdeleniya (zav. B.A. Tokareva) Gorodskoy detskoy
klinicheskoy bol'nitsy No.1, Moskva.
(CONJUNCTIVA---DISEASES) (CORTICOSTEROIDS)

TOKAREVA, V.A., kand.med.nauk; GEYMOS, Ye.K., vrach

Pemphigus and its treatment. Oft. zhur. 16 no.1:3-7 '61.

(MIRA 14:3)

1. Iz kafedry glaznykh bolezney (zav. - prof. N.A.Pletneva) II
Moskovskogo meditsinskogo instituta i 1-y detskoy klinicheskoy
bol'nitsy.

(PEMPHIGUS)

GEYN, E.E.

Device for recording the radiation pattern of antennas, operating
on medium frequencies. Vest. svyazi 24 no.1:9-10 Ja '64.
(MIRA 17:3)

1. Starshiy inzh. laboratorii TSentra tekhnicheskogo radiokontrolya.

GEYN, E.E.


Semigraphical method for calculating radio wave propagation
of the soil. Vest. svyazi 24 no.10:8-9 0 '64.

(MIRA 17:12)

1. Starshiy inzh. Tsentra tekhnicheskogo radiokontrolya.

"APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000515010011-7



APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000515010011-7"

GEYN, L. G.

5

1. Identification of cadmium by means of
 flame spectroscopy. ANALYTICAL PROCEDURE
 1.000 g. of sample is dissolved in 10 ml. of 10% NaOH.
 1.000 g. of 10% NaOH is added to the solution.
 1.000 g. of 10% NaOH is added to the solution.
 1.000 g. of 10% NaOH is added to the solution.
 The central spin containing only Cd is heated and saturated with
 NaOH. The central solution is treated
 dropwise with H₂O. A yellow ppt. of CdS appears.
 The min. amount of Cd detectable is 0.025 mg per
 ml. or 1.00 µg in one drop. C. S. Smith

PM

GEYN. L. G.

6

✓ Differential polarographic titration. N. Ya. Kulagin,
I. G. Chibrikov, and A. A. Babitskii (Pribor. 1981, 24(10)).
Zvezdnyy Znak 21, 1981, 1110-1111. The results of polarog-
raphy are applied to acid-base and oxidation-reduction
titrations are described. The setup for differential polarog-
raphy consists of 2 Pt wire electrodes connected in the
usual polarographic circuit, across which is connected an
external source of p.d. with circuit carrying a high resistance
(20 megohms). After initial balancing, the titration is run
conventionally, the parameters being believed to vary
after each addition of the reagent. The ratio of the opposed
voltages must be selected experimentally for each case. The
results of such titrations are curves with sharp and readily
determined breaks at the respective end points.
G. M. Kiselevich

CH

①

BE

KHLOPIN, N.Ya.; GAYN, L.G.

Behavior of the system of bromate-bromide-antipyrine under conditions of polarographic differential titration. Trudy Perm. farm. inst. no.1:134-137 '59. (MLWA 15:1)

1. Permskiy farmatsevticheskiy institut, kafedra neorganicheskoy i analiticheskoy khimii.
(ANTIPYRINE) (BROMINATION) (POLAROGRAPHY)

GEYN, L.G.

Detection of ions of bivalent and trivalent iron by means of phenol
and hydrogen peroxide. Trudy Perm. farm. inst. no.1:148-158 '59.
(MIRA 15:1)

1. Permskiy farmatsevticheskiy institut, kafedra neorganicheskoy i
analiticheskoy khimii.
(IRON-ANALYSIS)

USSR/ Analytical Chemistry - General Questions

G-1

Abs Jour : Referat Zhur - Khimiya, No4, 1957, 11982

Author : Lapin L.N., Geyn V.O.

Inst : Commission on Analytical Chemistry of the Academy of Sciences USSR

Title : Use of Basic Dyes of Diamino-Triphenylmethane Series for the Detection of Minute Amounts of Antimony, Gold and Thallium

Orig Pub : Tr. Komis. po analit. khimii. AN SSSR, 1956, 7(10), 217-222

Abstract : To decrease the effect of excess HCl and NaNO_2 on sensitivity of detection of Sb as SbCl_6^- by means of dyes of di- and triphenylmethane series, it is recommended to dissolve the resulting compound in organic solvents. Compound of Crystal Violet with SbCl_6^- ion passes readily into benzene and its homologues, imparting to them a blue-violet coloration. On using derivatives of diamino-triphenylmethane (Malachite Green and Brilliant Green) more reliable results are obtained. To 1 ml of the solution under study,

Card 1/2

USSR/ Analytical Chemistry - General Questions

G-1

Abs Jour : Referat Zhur - Khimiya, No 4, 1957, 11982

containing Sb as Sb^{3+} , are added 3 ml 5 N HCl, 2 drops 5% solution $NaNO_2$ and 0.2 ml of 0.5% solution of Brilliant Green in alcohol (diluted with water in the proportion 1:3). To the resulting orange-yellow solution is added 1 ml of toluene. In the presence of Sb the toluene acquires a green coloration the intensity of which increases with concentration of Sb. Detectable minimum is 0.05 % Sb. An analogous reaction with Brilliant Green is produced by $AuCl_3$ and $TlCl_3$, which interact as $AuCl_4^-$ and $TlCl_4^-$. Detectable minima are 0.5 % Au and 2 % Tl. Auramine makes it possible to detect Sb in the presence of Au and Tl, but the reaction is not sufficiently sensitive (detectable minimum 5 % Sb).

Card 2/2

GEYNAL, Ya. [Heinal, J.]; GRDLICHKA, Z. [Hrdlicka, Z.]; VRUBEL', I.

Protective effect of chlortetracycline on the vitality of exsanguinated tissues and organs. Antibiotiki 5 no.6:25-30 N-D '60.

(MIRA 14:3)

1. Institut klinicheskoy i eksperimental'noy khirurgii. Praga-Krch.
(AUREOMYCIN) (HEMORRHAGE)

GEYNBOM, L.

Strengthen the control of meal beetles infesting grain.
Muk.-elev. prom. 28 no.7:7 J1 '62. (MIRA 15:9)

1. Dnepropetrovskiy tekhnologicheskoy institut.
(Meal worms)

GEYNBOM, L.

Observing the temperature of stored corn. Muk.-elev. prom.
29 no.7:5-6 J1 '63. (MIRA 17:1)

1. Dnepropetrovskiy tekhnologicheskii tekhnikum.

GEYNE, A. G.

"The Utilization of the Interrelationships Between Stimulations for
Studying Patients Having Organic Disturbances of the Central Nervous System."
Sov Med Sci, Tbilisi State Medical Inst, Tbilisi, 1955. (KL, No 15, Apr 55)

SO: Sum. No. 704, 2 Nov 55 - Survey of Scientific and Technical Dissertations
Defended at USSR Higher Educational Institutions (16).

USSR/Human and Animal Physiology - The Nervous System.

T

Abs Jour : Ref Zhur Biol , No 3, 1959, 13243

Author : Popkhadze, N.A., Geyne, A.O.

Inst : -

Title : Investigation of Cortical Dynamics in Normal Individuals and Patients by Observation of Behavior Manifestations.

Orig Pub : Zh. vyssh. nervn. deyat-sti, 1957, 7, No 4, 501-509

Abstract : In patients with intero-neurasthenic syndrome, who had been blind-folded, the conditioned reflex (CR) was developed by means of pressure of the leg on a pedal in response to verbal instruction when at first a wooden ball with a diameter of 45 cm was placed in the hand and then a differentiation with a ball of 42 mm. The stress stimulus (SS) - a ball of 100 mm - produced CR, but when it was followed by the ball of 45 cm CR did not occur. After 12 - 25 minutes the

Card 1/2

USSR/Human and Animal Physiology - The Nervous System.

T

Abs Jour : Ref Zhur Biol., No 3, 1959, 13243

reactions were restored (normally after 2 - 3 minutes).
SS - ball of 18 mm - did not evoke CR, but the ball
of 42 mm 2 - 3 minutes after SS provoked CR (normally -
immediately after SS). Then 2 balls were presented at
the same time -- 40 mm to the right and 66 mm to the
left, and the subjects were requested to press the
pedal on the side of the larger ball. After a number
of combinations 2 balls of 40 mm were presented.
With this CR appeared on the right in some of the sub-
jects. 2 - 5 minutes after SS - the ball of 18 mm -
presentation of 2 balls of 40 mm induced CR on the
left. The indices were normalized as a result of the
therapy. -- M.I. Lisina

Card 2/2

- 109 -

GEYNE, A.O.,

USSR / Pharmacology, Toxicology, Narcotics and Hypnotics

U-2

Abs Jour : Ref. Zh.-Biol., No 2, 1958, No 7924

Author : Mindadze, A.A., Geyne, A.O.

Inst :

Title : The Effect of Small Doses of Alcohol on the Cortex of the Cerebral Hemispheres

Orig Pub : Tr. Tbilissk. in-t, 1957, 11, 91-100

Abstract : The relationship between conditioned stimuli was studied in 10 healthy subjects before and following the drinking of 50 g of 40% alcohol. Prior to this a conditioned motor reflex was developed to a cutaneous kinesthetic stimulation caused by placing a sphere with a 67 mm diameter in the subject's palm and its differentiation from a sphere with a 40

Card : 1/2

С-15 YAF 11:1

POPKHADZE, N.A.; GRYNE, A.O.

Method for detecting disorders in kinesthetic sensitivity of the muscles and joints. Zhur.nevr. i psikh. Supplement:47 '57.
(MIRA 11:1)

1. Klinika nervnykh bolezney pediatricheskogo i sanitarno-gigiyenicheskogo fakul'tetov Tbilisskogo meditsinskogo instituta.
(CONDITIONED RESPONSE) (MUSCULAR SENSE)

GEYNE, G.A.

Blocking of the vegetative ganglia and trunks of the extracardial nerve apparatus of the heart in acute coronary insufficiency under experimental conditions. Grud.khir. no.3:42-49 '61.

(MIRA 14:9)

1. Iz kafedry operativnoy khirurgii i topograficheskoy anatomii (zav. - zasluzhennyy deyatel' nauki prof. V.V. Babuk) Minskogo meditsinskogo instituta.

(CORONARY HEART DISEASE) (AUTONOMIC DRUGS)

GEYNE, G.A. [Heine, H.A.]

Effect of novocaine block of the left stellate ganglion on morphological changes in the myocardium following ligation of the left coronary artery at a high level. Vestsi AN BSSR. Ser. biol. nav. no.3:54-59 '61. (MIRA 14:10)

(NERVOUS SYSTEM, SYMPATHETIC)
(HEART...MUSCLE) (CORONARY VESSELS)

GEYNE, G.A.

Normalization of cardiac rhythm under the influence of a novocaine block of the stellate ganglion in experimentally induced acute coronary circulatory disorders, Zdrav.Belor. 5 no.12:29-32 D '59.

(MIRA 13:4)

1. Kafedra operativnoy khirurgii topograficheskoy anatomii (zav. - prof. V.V. Babuk) Minskogo meditsinskogo instituta.

(NOVOCAINE) (CORONARY VESSELS--DISEASES) (HEART)

GEYNE, G. A. Cand Med Sci -- "Acute experimental coronary ~~deficiency~~ ^{deficiency} and
certain methods of controlling it." Smolensk, 1961 (Smolensk State Med Inst).
(KL, 4-61, 207)

-329-

GEYNE, G.A.

Effect of a bilateral cervical vagosympathetic novocaine block on morphological changes in the myocardium after a high ligation of the left coronary artery. Zdrav. Bel. 9 no.2:21-24 F'63. (MIRA 16:7)

1. Kafedra operativnoy khirurgii i topograficheskoy anatomii (zaveduyushchiy - zasluzhennyy deyatel' nauki prof. V.V.Babuk) Minskogo meditsinskogo instituta.
(NOVOCAINE) (VAGUS NERVE) (CORONARY VESSELS—LIGATURE)
(HEART—MUSCLE)

GEYNE, G.A. [Heine, H.A.]

Restoration of coronary blood circulation during the acute period
of experimental myocardial infarct. Vestsi Akad. BSSR. Ser. biol.
nav. no. 4:74-79 '63. (MIRA 17:8)

GUDKOVA, Ye.I., MIRONOVA, K.A., KUZ'MINSKIY, A.S., GEYNE, G.O.

Second outbreak of listerellal anginas in a populated area.
Zhur.mikrobiol. epid. i immun. 29 no.9:24-28 S'58 (MIRA 11:10)

1. Iz Instituta ukha, gorla, i nosa.
(TONSILLITIS, epidemiology,
in Russia, caused by Listeria (Rus))
(LISTERIA, infections,
tonsillitis, epidemiol. in Russia (Rus))

GEYNE, V.


Work experience of the propaganda section. Kryl.rod. 3 no.4:8
Ap '52. (MIRA 8:8)

1. Chlen Ismail'skogo oblastnogo orgkomiteta Dobrovol'nogo
obshchestva nodaystviya armii, aviatsii i flotu.
(Military education)

GEYNE, V.

Set up clubs for primary organizations. Voen.znan.31 no.4:8 Ap'55.
(MLRA 8:10)

1. Chlen oblastnogo komiteta Dobrovol'nogo obshchestva sodeystviya
armii, aviatsii i flotu
(Kuybyshev Province--Military education)



GEYNE, Ye. I

May 49

USSR/Chemistry - Oxidation
Cellulose

"The Chemistry of Viscose and Xanthogenates: III, Oxidation Reactions in Alkalicellulose and Viscose," S. N. Danilov, N. M. Grad, Yel I. Geyne, Lab of Chem Processing of Cellulose, Leningrad Technol Inst in Lensovet, 17 pp

Zhur Obshch Khim, Vol 19, No 5, p 826

Study of the oxidation of alkalicellulose in the presence of retarding and accelerating impurities showed that sodium sulfite alone or with sodium sulfide markedly retarded the activity while the sulfide alone produced a less marked effect. The retardation in this case was actually preceded by an acceleration in the first moment of the reaction. Noted similar retarding effects for sodium sulfite and sodium sulfide in viscose. Demonstrated that the xanthogenates of ethyl alcohol, diacetone glucose, and cellulose react with oxygen to form dixanthogenides. Submitted 18 Jan 48.

PA 67/49T58

GLOTOV, V.N.; GEYNE, Ye.I.

Method of preparing coruleum, a pigment for art painting.
Lakokras.mat. i ikh prim. no.2:18-20 '61. (MIRA 14:4)
(Pigments)

13493-66 (N) INT(m)/INT(j)/T RM
 ACC NO: AF6001681 SOURCE CODE: UR/0303/65/000/006/0025/0027
 AUTHORS: Gurevich, Ye. S.; Glotov, V. N. (deceased); Gayne, Ye. I. 436
 ORG: none B
 TITLE: Kinetics of leaching of poisons from coatings of antifouling paints 6,445
 SOURCE: Lakokrasochnyye materialy i ikh primeneniye, no. 6, 1965, 25-27
 TOPIC TAGS: vinyl, protective coating, pigment, copper compound, sea water/ KhV 53
 perchlorovinyl resin based paint, KhC 79 chlorovinyl and vinyl acetate copolymer
 based paint, A 15 vinyl acetate
 ABSTRACT: The effect of mineral, organic, and chelate additives upon leaching of
 copper from coatings of antifouling paints was investigated. The work was undertaken
 as an expansion of previous investigations by the authors (Lakokrasochnyye materialy i
ikh primeneniye, No. 6, 53(1964); V. N. Glotov. Zav. lab., 30, No. 1, 111, 1964) in
 order to devise new and more economical antifouling coatings than those containing
 the scarce and expensive cuprous oxide. Rates of leaching of copper as the poisonous
 material from various types of antifouling coatings as functions of time are
 illustrated in Fig. 1. The investigated paints were of type KhV-53, perchlorovinyl
 resin based, and KhC-79, based on a copolymer of chlorovinyl with vinyl acetate A-15.
 The controls contained cuprous oxide as the only pigment and poison. Experimental
 work and testing at the Black Sea have shown that most of the chelating compounds
 Cord 1/2 UDC: 667.613.3:620.193.23

L 13493-66
ACC NO: AF6001681




Fig. 1. Rate of leaching of poisons as function of time for antifouling paints; curves 1, 2 - soluble film-forming; curves 3, 4 - contact type; 1, 3 - theoretical rate; 2, 4 - practical rate.

accelerated the leachability of copper poison, lowering the consumption of copper, increasing the effectiveness of antifouling coating, and extending the period of service of the paint. L. I. Shcherbakova, A. I. Smirnova, N. V. Skoropostishnaya, and A. I. Nagel' participated in this work. This study was a development of work by M. A. Dolgopolskaya and B. S. Gurevich (Trudy Sevastopol'skoy biologicheskoy stantsii AN UkrSSR, t. 14, t. 13, t. 11, 1959) who first investigated antifouling paints with the application of chelating compounds (derivatives or salts of 8-oxyquinoline). Orig. art. has: 3 tables and 1 figure.

SUB CODE: 11, 07/

SUBM DATE: none/

ORIG REF: 006/

OTH REF: 003

Card 2/2

GEYNEMAN, A.V.; MEDZHIBOZHSKIY, M.Ya.

Effect of temperature on the speed of the decarburization
process in open-hearth furnaces. Izv. vys. ucheb. zav.;
chern. met. 6 no.8:57-59 '63. (MIRA 16:11)

1. Sibirskiy metallurgicheskiy institut.

MEDZHIFOMSKIY, M.Ya., doktor tekhn. nauk; KURAPIN, B.S.; GAYNEMAN, A.V.;
IVORYANINOV, V.A.; MOISEYENKO, A.I.; LOSHCHEV, V.Ya.

Nitrogen-content in the metal during the blowing of the
open-hearth furnace bath with compressed air. Met. 1
gornorud. prom. no.6:23-26 N-D '65. (MIRA 18:12)

DAGAYEV, V.N.; GREYNEMAN, N.V.

A case of an aneurysm of the abdominal aorta into the duodenum.
Khirurgiia 32 no.6:72-73 Je '56. (MLRA 9:10)

1. Iz Tul'skoy gorodskoy bol'nitsy imeni N.A.Semashko (glavnyy
vrach i zav. khirurgicheskim otdeleniyem - doktor meditsinskikh
nauk V.F.Dagayev)

(AORTIC ANEURYSM, case reports rupture
abdom., into duodenum)

S/148/60/000/006/012/016/XX
A161/A030

AUTHORS: Medzhibozhskiy, M.Ya.; Zinov'yev, V.T.; Geyneman, A.V.

TITLE: The Effect of Some Factors on the Carbon Burning Rate in the Open-Hearth Furnace Bath

PERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy. Chernaya metallurgiya, 1960, No. 6, pp. 47 - 53

TEXT: Most authors consider that the oxidation of carbon in the open-hearth bath is limited by diffusion links, but some point out that it may also be limited by heterogeneous chemical reactions on the boundaries gas-slag, slag-metal and metal-bubbles. The question of in what region the process takes place, diffusion (limitation by diffusion links) or kinetic (chemical links) can be answered by determining the effect of the temperature on the speed of the process. With relatively slow chemical reactions the process rate will change with temperature in accordance with the Arrhenius equation

$$K = K_0 e^{-\frac{E}{RT}}$$

where K is the constant of the chemical reaction rate; K_0 the so-called preexpo-

Card 1/7

S/148/60/000/006/012/016/XX
A161/A030

The Effect of Some Factors on the Carbon Burning Rate in the Open-Hearth Furnace Bath

ponential factor; e the base of natural logarithms; E the activation energy, cal/mol; R the universal gas constant, 1.986 cal/mol °K; T the absolute temperature, °K. It follows that

$$E = \frac{4.575 \lg \frac{K_2}{K_1}}{\frac{1}{T_1} - \frac{1}{T_2}}, \quad (2)$$

K_1 and K_2 are constants of the chemical reaction speed at T_1 and T_2 temperatures. The process is in the kinetic region if the activation energy value is high ($> 50,000$ cal/mol), and in the diffusion region if it is relatively low ($> 30,000$ cal/mol). The authors studied over 400 heats in 380 and 190-ton open-hearth furnaces firing mixed coke and generator gas; temperature was measured by means of a submerged platinum-rhodium-platinum thermocouple. Over 1,200 carbon oxidation rate values were determined (v_C) at different temperatures (t_M) and C content in metal. The relations shown were verified by mathematical statistic methods. The effect of the C content in metal, of the specific surface area of the slag-metal boundary, and the quantity of oxygen was also studied. The results are illustrat-

Card 2/7

S/148/60/000/006/012/016/XX
A161/A030

The Effect of Some Factors on the Carbon Burning Rate in the Open-Hearth Furnace Bath

ed by diagrams (Figs. 1, 2, 3). The low activation energy proves that the process is in the diffusion region. Low activation energy values in the steel decarbonization process had also been obtained in laboratory experiments by S.I. Filippov (Ref. 1) and in shop experiments by P. Vallet (Ref. 7, Iron and Steel, 1955, No. 11). Conclusions: 1) The carbon burning rate does not depend on the carbon content in metal if $\%C > 0.3\%$. 2) The effect of the temperature of the metal on the carbon burning rate is not strong at superheating above the liquidus point ($> 20^\circ$) and corresponds with the apparent activation energy of the decarbonization process amounting to 15,000 - 22,000 cal/mol. This means that the process takes place in the diffusion range. 3) The decarbonization rate depends on the intensity of oxygen feed into the bath and is not limited by any of the heterogeneous reactions on the boundary's gas-slag, slag-metal and metal-bubbles. 4) At $\%C < 0.3\%$, the relation between v_C and $\%C$ is caused not by the fact that the decarbonization process is in the kinetic region, but by other causes that had been pointed out in the works (Refs. 1, 8, 9, 10). 5) The indirect effect of $\%C$ and t_M on v_C is very considerable with large additions of iron ore into the bath. The intensity of carbon burning through ore additions (which reduce the

Card 3/7

S/148/60/000/006/012/016/XX
A161/A030

✓
-


The Effect of Some Factors on the Carbon Burning Rate in the Open-Hearth Furnace Bath

bath temperature) is limited by the necessity to maintain the minimum superheating of metal. There are 3 figures and 12 references: 11 Soviet and 1 English.

ASSOCIATION: Sibirskiy metallurgicheskiy institut (Siberian Metallurgical Institute)

SUBMITTED: October 27, 1959

Card 4/7



S/148/60/000/006/012/016/XX
A161/A030

The Effect of Some Factors on the Carbon Burning Rate in the Open-Hearth Furnace Bath

Figure 1: The effect of C content on the C burning rate during the pure boiling period. (The figures at the points indicate the number of heats). a - boiling without ore in the 1,570 - 1,620° range (in 190-ton furnace) and in 1,540 - 1,580° (380-ton furnace); b - by the data from all samples taken during pure boiling. 1 - 190-ton furnace; 2 - 380-ton furnace.

Card 5/7

S/148/60/000/006/012/016/XX
A161/A030

The Effect of Some Factors on the Carbon Burning Rate in the Open-Hearth Furnace Bath




Figure 2: The effect of iron ore additions on the carbon burning rate at the beginning of the pure boiling time: 1 - 190-ton furnace; 2 - 380-ton furnace. (Ore consumption from 0.2 to 1.4 %/h).

Card 6/7

S/148/60/000/006/012/016/XX
A161/A030

The Effect of Some Factors on the Carbon Burning Rate in the Open-Hearth Furnace Bath




Figure 3: The effect of metal temperature on the carbon burning rate during the pure boiling time: 1 - the beginning of pure boiling time (ore consumption about 0.5%/h, or 0.17%, during the first 20 min); 2 - the beginning and the mid of the pure boiling time (without ore additions); 3 - the last 15 - 20 min of pure boiling.

Card 7/7

GETNIN, N.G., inzh.

Equivalent zero-sequence substitution circuits for several
closely running lines. Elektrichestvo no.8:28-32 Ag '60.

(MIRA 13:8)

(Electric lines--Overhead)

GEYNIN, N.G., inzh. (Leningrad)

Reactance of the mutual inductance of lines having different
potentials in zero-sequence equivalent circuits. Elektrichestvo
no.7:79-81 J1 '61. (MIRA 14:9)
(Electric power distribution)
(Equivalent circuits)

GEYNISMAN, G.V., inzh.; FEDOTOV, Ya.V., inzh.

Stability of polymetal rods. Sudostroenie 25 no.4:13-15 Ap '59.
(MIRA 12:6)

(Elastic rods and wires)

GEYNISMAN, N.V.

Morphological changes in influenzal encephalitis. Vop.psikh.1
nevr. no.7:240-248 '61. (MIRA 15:8)

1. Iz 2-y Leningradskoy psikhonevrologicheskoy bol'nitsy (glavnyy
vrach T.I.Nikolayeva, nauchnyy rukovoditel' - starshiy nauchnyy
sotrudnik instituta imeni V.M.Bekhtereva, A.N.Ageyeva).
(ENCEPHALITIS) (INFLUENZA)

GEYNISMAN, N.V.; SKOBLO, M.S.

Cases of unusual manifestations of brain tumors. Vop.psikh.i nerv.
8:297-304 '62. (MIRA 17:4)

1. Iz 2-y psikhonevrologicheskoy bol'nitsy (glavnyy vrach
T.I.Nikolayeva), Leningrad.

GEYNISMAN, YA. I.

Geynisman, Ya. I. "X-ray diagnosis of the later results of firearms injuries to the skull and brain", In the collection: Nevrologiya voyen. vrachev, Vol. 1, Moscow, 1949, p. 170-89.

SB: U-11, 17 July 1951, (Letopis 'Chernol'nykh Stroy, no. 1, 1949)

SETHLEMAN, Ya. I.

Ya. I. Sethleman, P. O. Trenchell, and P. I. Iosadovich. "The results of X-ray therapy of some results of fibrous injuries to the brain", in the collection: Sanitarnaya voyennaya vremeni, Vol. I, Moscow, 1943, p. 273-92.

SO: U-411, 17 July 1953, (Letopis "Zhurnal 'nykh Statey, no. 20, 19.9)

СЫНОВИЯ", №. 1.

1921. К вопросу о физических рентгеновских лучах и о их
о нестрел'ных повреждениях, повреждениях. ... : ...
1., 1949, с. 63-64.

80: "Летопись" ... , "ол. 10, Москва, 1949

GEYNISMAN, Ye. I.; ZHIRMUNSKAYA, Ye. A.

Mechanism of action of roentgen rays on reflex excitability. *Fiziol.*
zh. SSSR 38 no.3:312-318 May-June 1952. (GLML 23:2)

1. Institute of Neurology, Academy of Medical Sciences USSR.

GEYNISMAN, Ya.I., professor; REYNBERG, S.A., redaktor.

[Delayed and directed pneumoencephalography; importance of this method of examination in sequelae of brain trauma] Zamedlennaiia i napravlennaiia pnevmontsefalografii; znachenie etogo metoda issledovaniia pri posledstviakh travmaticheskikh povrezhdenii golovnogo mozga. Moskva, Medgiz, 1953. 209 p. (MLRA 7:4)
(Encephalography)

GETHISMAN, Ya. I.; ZHIRMUNSKAYA, Ye. A.

~~SECRET~~
Mechanism of action of roentgen rays on the functional
state of the central nervous system; electroencephalo-
graphic data. Vest. rentg., Moskva No.2:5-15 Mar-Apr.
1953 (CML 25:5)

1. Of the Institute of Neurology (Director -- Prof. N.V.
Konevalov, Active Member AMS USSR) of the Academy of Medical
Sciences USSR.

GEYNISMAN, Ya.I., professor; ARUTYUNOV, A.I., professor, direktor.

Pneumoencephalographic methods. Vest.rent.i rad. no.3:63-67 My-Je '53.
(MLRA 6:8)

1. Nauchno-issledovatel'skiy institut neyrokhirurgii Min'isterstva zdra-
vookhraneniya USSR. (Encephalography)

GEYNISMAN, Ya.I., professor

Significance of tomography in diagnosis of arachnoid endotheliomas
of the olfactory region. Vop. neirokhir. 18 no.3:7-14 My-Je '54.
(MLRA 7:8)

1. Iz Nauchno-issledovatel'skogo instituta neyrokhirurgii Mini-
sterstva zdavookhraneniya USSR.

(NASAL CAVITY, neoplasms,

*diag., tomography, olfactory region)

GEYNISMAN, Ya.I., professor

Significance of sectional craniography in diagnosis of arachno-
endothelimas (Meningiomas) of the olfactory region. Vop. neirokhir.
18 no.4:3-8 J1-Ag '54. (MLRA 7:10)

1. Iz Nauchno-issledovatel'skogo instituta neyrokhirurgii Minister-
stva zdoravookhraneniya USSR.

(FRONTAL LOBE, neoplasms,

*meningioma of olfactory region, diag., tomography)

(NOSAL CAVITY, neoplasms,

*meningioma of olfactory region, diag., tomography)

(MENINGIOMA,

*olfactory region, diag., tomography)

GEYNISMAN, Ya.I., prof.; SOROCHINSKIY, TS.M.; DANILENKO, G.S.

Craniography in the diagnosis of brain tumors. Vrach.delo no.8:809-
813 Ag '59. (MIRA 12:12)

1. Otdel neyrorentgenologii (zav. - prof. Ya.I. Geynisman) Ukrainskogo
instituta neyrokhirurgii.
(BRAIN--TUMORS) (SKULL--RADIOGRAPHY)

GEYNISMAN, Ya.I., prof.; SOROGHINSKIY, TS.M.; TANANAYKO, P.G.

Delayed and controlled pneumoencephalography in subtentorial tumors [with summary in English, p.54]. Vop.neirokhir. 22 no.6:3-7 N-D '58. (MIRA 12:2)

1. Ukrainskiy nauchno-issledovatel'skiy institut neyrokhirurgii.
(BRAIN NEOPLASMS, diagnosis,
subtentorial, pneumo-encephalography (Rus))

GEYNISMAN, Ya.I., prof.

"X-ray diagnosis of neurological disorders in children" by N.N.
Alt'gauzen. Reviewed by IA.I. Geinisman. Vop.neirokhir. 23 no.1:59
'59. (MIRA 12:3)

(NERVOUS SYSTEM--DISEASES)
(DIAGNOSIS, RADIOSCOPIC)
(CHILDREN--DISEASES)

GEYNISMAN, Ya.I., prof. (Kiyev, ul. Vorovskogo, d.27, kv.7)

Tomography in the diagnosis of meningiomas of the alae of the basilar bone. Vest.rent.i rad. 34 no.2:3-8 Mr-Apr '59. (MIRA 13:4)

1. Iz neyrorentgenologicheskogo otdela (nauchnyy rukovoditel' - prof. Ya.I. Geynisman) Ukrainskogo nauchno-issledovatel'skogo instituta neyrokhirurgii (direktor - zasluzhennyy deyatel' nauki prof. A.I. Arutyunov).
(MENINGIOMA, diagnosis,
tomography in meningioma of basilar bone
alae (Rus))

GEYNISMAN, Ya.I.; PANKEYEVA, L.P.; PCDOL'SKIY, F.D.

Radiographic changes in the skull in brain tumors of varying
histostructure. Probl.neirokhir. 4:131-147 '59. (MIRA 13:11)
(SKULL--RADIOGRAPHY)
(BRAIN--TUMORS)

GEYNISMAN, Yu.Ya.

Dehiscence of cranial sutures in intracranial hypertension [with
summary in English, p.64] Vopr.neirokhir. 22 no.4:49-53 J1-Ag '58
(MIRA 11:9)

1. Kafedra patologicheskoy anatomii I Moskovskogo ordena Lenina
meditsinskogo instituta imeni I.M. Sechenova.

(BRAIN, NEOPLASMS, compl.

intracranial hypertension with dehiscence of cranial
sutures (Rus))

(CRANIUM,

suture dehiscence in intracranial hypertension
caused by tumors (Rus))

GEYNISHAN, Yu.Ya.; GOL'TS, H.V. (Moskva)

Experimental comparison of methods for the intramedullary fixation of fresh fractures. Eksp.khir. 4 no.3:47 My-Je '59.

(MIRA 12:8)

(FRACTURES)

GEYNISMAN, Yu.Ya.

Mechanism of the action of aminazine and imizine. Dokl. AN SSSR
141 no.2:461-463 N '61. (MIRA 14:11)

1. Institut vysshey nervnoy deyatel'nosti i neyrofiziologii AN SSSR.
Predstavleno akademikom V.N.Chernigovskim.
(CHLORPROMAZINE) (IMIPRAMINE) (BRAIN)

GEYNISMAN, Yu.Ya.

Histochemical changes in the reticular formation of the brain
stem of animals under the influence of aminazine and imizine.
Trudy Inst. vys. nerv. deiat. Ser. patofiziol. no.9:168-173 '61.
(MIA 15:4)

(BRAIN) (CHLORPROPAZINE) (IMIPRAMINE)

39201
S/246/62/062/002/004/006
1015/1215

AUTHOR: Geynisman, Yu. Ya.

TITLE: Effect of aminazin and imizin on morphological changes in the brain of animals

PERIODICAL: Zhurnal nevropatologii i psikiatrii imeni S. S. Korsakov, v. 62, no. 2, 1962, 190-196

TEXT: According to the author no other comparative studies on the morphologic changes in CNS following aminazin and imizin administration had been done. The experiments were carried out on 176 male, sexually mature, albino rats, weighing 170-250 g. Three series of experiments were carried out: 1) a single introduction of 5 mg/kg b.w. of aminazin and imizin; 2) a single introduction of 20 mg/kg of these drugs, and 3) a chronic experiment with 5 mg/kg aminazin and imizin which were administered once a day for 30 days. Further data on timing and staining methods for histologic examination are given. The main histologic findings were: no changes in myelinated fibres, hyperchromasia of cortical neurons in the II, III and V layers and an increase in argyrophil granules, proteins, RNA, and SH groups in these cells. These changes were observed soon after the administration of the drugs. Hypochromasia and vacuolisation of cells in the reticular formation as well as a marked glial reaction which differed with the time intervals since the administration of the drugs, were significant later on. Thus, a biphasic effect of aminazin and imizin was established. This was shown also in physiological investigations. The second phase was found to begin earlier after the introduction of imizin than after aminazin. There are 3 figures.

Card 1/2

GEYNISMAN, Yu.Ya.

Comparative analysis of morphological changes in the cortex
and subcortical formations in the animal brain under the
action of aminazine and imizin on healthy animals. Trudy Inst.
vys.nerv.deiat. Ser.fiziol. 7:288-299 '62. (MIRA 16:2)
(IMIPRAMIN) (CHLORPROMAZINE) (BRAIN)

ALEKSEICHENKO, M.M.; GENTSEMAN, N.Ya.; GUSEV, V.G.

Morphophysiological data on the analysis of the mechanism of
the effect of aziridine on the central nervous system. *Sov.
Physiol. J.* 14 no. 4/1970 p. 100-104.

(MIRA 17:12)

L. Laboratory of Morphology of the Central Nervous System and
Cardio and Reflexes, Institute of Brain, Nervous Activity and
Neurophysiology, U.S.S.R. Academy of Sciences, Moscow.

100-443887-100

08/09/06 09:09 AM

THE UNIVERSITY OF CHICAGO PRESS

3.2.2.2. *Effect of the temperature on the adsorption*

84-681

THE FORD CREDIT CORPORATION
10000 FORD DRIVE
DEARBORN, MICHIGAN 48120

For 40 min in 15 sec and 10 sec, 50% of the

GETNISMAN, Yul'ya. (Moskva)

Shifts in the content and concentration of cytoplasmic ribonucleo-
proteins in functional changes in the neurons. Arkh. nat. 27 (MIRA 18:10)
no.10:76-78 '65.

1. laboratoriya morfologii tsentral'noy nervnoy sistemy (zav. -
prof. M.M.Aleksandrovskaya) Instituta vysshey nervnoy deyatelnosti
i nevrofiziologii (direktor - chlen-korrespondent AN SSSR prof. I.I.
Asratyan) AN SSSR.

L 1596-66

ACCESSION NR: AF5024775

UR/0219/64/058/009/0080/0086

AUTHOR: Aleksandrovskaia, M. N.; Gaynisman, Yu. Ya.; Samoylova, L. G. 208

TITLE: Structural and metabolic changes of the brain of animals with disturbances of higher nervous activity following repeated administration of chlorpromazine

SOURCE: Byulleten' eksperimental'noy biologii i meditsiny, v. 58, no. 9, 1964, 80-86

TOPIC TAGS: medical experiment, rat, brain, central nervous system, chlorpromazine, biologic metabolism, neurology

ABSTRACT: Thirty-five male white rats received daily injections of 5 milligrams/kilogram of chlorpromazine for a period of 30 days. During the first 3 weeks there was a depression of the conditioned reflex activity and marked histological and histochemical changes in the cerebral cortex, the diencephalon nuclei, and stem reticular formations. By the 30th day, the disturbances of higher nervous activity and the microscopic brain lesions became less pronounced. For 3 weeks after injections were stopped there was increased excitation of the higher parts of the central nervous system, and

Card 1/2

L 1596-66

ACCESSION NR: AP5024775

0

the brain's structural and metabolic changes acquired a different character, being chiefly localized in the cortical neurons. On the 60th day, the animals' behavior, conditioned reflex activity, and state of cerebral tissue did not differ from those of the controls. It is concluded that the functional, metabolic and structural changes of the higher parts of the central nervous system of animals caused by repeated administration of chlorpromazine are reversible. Orig. art. has: 12 figures, 1 table.

ASSOCIATION: Laboratoriya morfologii tsentral'noy nervnoy sistemy, Instituta vysshey nervnoy deyatel'nosti i neyrofizologii, AN SSSR. Moscow (Laboratory of the Morphology of the Central Nervous System, Institute of Higher Nervous Activity and Neurophysiology); Laboratoriya uslovykh refleksov, Instituta vysshey nervnoy deyatel'nosti i neyrofizologii, AN SSSR. Moscow (Laboratory of Conditioned Reflexes, Institute of Higher Nervous Activity and Neurophysiology)

SUBMITTED: 08Jul63

ENCL: 00

SUB CODE: LS

NR REF SOV: 010

OTHER: 013

JPRS

Card 2/2 *SP*

ALEKSANDROVSKAYA, M.M.; GEYNISMAN, Yu.Ya.; MATS, V.N.

Glial-neuronal relations during intensified functioning of the neurons according to morphological research data. Zhur. nevr. i psikh. 65 no.2:161-167 '65. (MIRA 18:9)

1. laboratoriya morfologii TSentral'noy nervnoy sistemy (zaveduyushchiy - prof. M.M. Aleksandrovskaya) Instituta vysshey nervnoy deyatel'nosti i neyrofiziologii (direktor - prof. E.A. Asratyan) AN SSSR, Moskva.

GHYNTS, V.V.

Transition of epidemic hepatitis to chronic form. *Sov. med.* 28
no.5:90-93. May '65. (MIRA 12:5)

1. Klinicheskoye otdeleniye Tadjikskoye Institut krayevoy
meditsiny (dir. - chlen-korrespondent AN Tadjikskoy SSR Kd.Kn.
Mamurov) AEN SSR i Gorodskoye infektatsionnyy tsentr (glavnyy
vrach A.V.Markova), Dushanbe.

GEYNOVA, L.A.; MATOVSKIY, I.M.; ODINTSOVA, K.P.; SHAPIRO, A.Sh.
(Chelyabinsk)

Prophylaxis for angina incidence at industrial enterprises in
Chelyabinsk. Zdrav. Ros. Feder. 7 no.8:30-32 Ag'63.
(MIRA 16:10)

(CHELYABINSK — TONSILS — DISEASES)
(CHELYABINSK — INDUSTRIAL HYGIENE)

3 EYK 15 11 11 4
ZELIKMAN, A.L.; GMYNRIKH, A.K.

Fertility of cyclops in culture media of various density. Dokl.
AN SSSR 60 no.5:925-927 My '48. (MLRA 10:8)

1. Institut zoologii Moskovskogo gosudarstvennogo universiteta
im. M.V. Lomonosova, Predstavleno akademikom I.I. Shmal'gausenom.
(Copepoda)

GEYNRIKH, A. K.

GEYNRIKH, A. K.: "Seasonal phenomena among the zooplankton of the Bering Sea". Moscow, 1955. Acad Sci USSR, Inst of Oceanology. (Dissertation for the Degree of Candidate of Biological Sciences)

SC: Knizhnaya Letopis', No. 40, 1 Oct 55

GEYRIKH, A.K.

Seasonal races in *Calanus tonus* in the Bering Sea. Dokl. AN SSSR 109
no.2:403-405 J1 '56. (MIRA 9:10)

1. Predstavleno akademikom N.N. Pavlovskim.
(Bering Sea—Copepoda)

GEYNRIKH, A.K.

Biometric data on Chaetognaths and their propagation in the
western regions of the Bering Sea. Dokl. AN SSSR 110 no.6:
1105-1107 0 '56. (MLRA 10'2)

1. Predstavleno akademikom Ye.N. Pavlovskim.
(Bering Sea--Nemathelminthes)

GEYNRIKH, A.K.

Copepoda production in the Bering Sea. Dokl.AN SSSR 111 no.1:199-201
M-D '56. (MLRA 10:2)

1. Predstavleno akademikom Ye.N.Pavlovskim.
(Bering Sea--Copepoda)